

CHEMICAL WEED CONTROL FOR WHEAT AND BARLEY IN THE LOW DESERTS OF ARIZONA

Barry Tickes, University of Arizona

Herbicide	Rate (lb ai/A)	Grasses				Mustard			Chenopodium			Composite		Other		Comments
		Canarygrass	Wildoat	Rabbitfoot Grass	Italian Ryegrass	London Rocket	Shepardspurse	Wild Mustard	Nettleleaf Goosefoot	Lambsquarter	Russian Thistle	Annual Sowthistle	Prickly Lettuce	Silversheath Knotweed	Little Mallow	
Banvel 4S & Clarity 4S	0.06 to 0.12	○	○	○	○	●	●	●	●	●	●	◐	●	◐	Less crop safety than MCPA, 2,4-D & Buctril especially on barley. Should be applied prior to 5 leaf stage of growth. <u>Drift Hazard</u> . Tank mixes with other herbicides may increase potential crop injury.	
2,4-D	0.25 to 0.50	○	○	○	○	●	●	●	●	●	●	●	●	◐	◐	<u>Drift Hazard</u> to a wide range of broadleaf crops. Should be applied after tillering & before boot stage of growth. Tank mixes with other herbicides may increase potential of crop injury.
MCPA	0.25 to 1.5	○	○	○	○	●	●	●	●	●	◐	●	◐	◐	<u>Drift Hazard</u> but safer to cotton & alfalfa than 2,4-D. Apply after tillering & before boot stage. Tank mixes will increase potential crop injury.	
Buctril 2E	0.25 to 0.375	○	○	○	○	●	●	●	●	●	●	●	●	○	○	This is a contact herbicide & will only control small weeds. Less drift hazard. Can be applied from emergence to boot stage of crop development.
Aim 2E	0.008 to 0.016	○	○	○	○	○	◐	●	◐	◐	◐	◐	○	◐	●	Contact herbicide with little or no residual activity. Good coverage to actively growing weeds essential.
Achieve 40DG	0.18 to 0.24	●	●	*	●	○	○	○	○	○	○	○	○	○	○	Best weed control will result if applied after most of the weeds have emerged and the largest ones are no more than 4 to 7 leaves.
Hoelon 3E	0.75 to 1.0	◐	●	○	●	○	○	○	○	○	○	○	○	○	○	Canarygrass with 3 or more leaves will not be controlled. Crop oil will improve control but may increase temporary crop injury.
Avenge 2S	0.625 to 1.0	○	●	○	○	○	○	○	○	○	○	○	○	○	○	May cause unacceptable crop injury to durum & other wheat varieties. Will control wild oat up to 5 leaf stage. Resistance of wild oats has been noted in several Western states.
Stinger 3E	0.1 to 0.125	○	○	○	○	○	○	○	○	○	◐	●	●	○	○	Avoid drift to sensitive crops. Apply from 3 leaf to boot stage of crop.

● Good Control ◐ Partial Control ○ No Control * Too little local data to evaluate.

Any products, services or organizations that are mentioned, shown or indirectly implied do not imply endorsement by the University of Arizona. Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, James A. Christenson, Director, Cooperative Extension, College of Agriculture and Life Sciences, The University of Arizona. The University of Arizona is an equal opportunity, affirmative action institution. The University does not discriminate on the basis of race, color, religion, sex, national origin, age, disability, veteran status, or sexual orientation in its programs and activities.