Lepidopterous Larvae Management in Desert Produce Crops, 2016



	IRAC 1	Beet	Cabbage	Corn	THE UNIVERSITY OF ARIZONA COOPERATIVE EXTENSION Yuma Agricultural Center
Product	MOA	armyworm	looper	earworm	Comments*
Lannate	1A	•••		•••	Tank mix with another product for broad spectrum Lep activity; provides thrips control;
Laillate	14		· ·	***	PHI: 10 d on lettuce; Use rates above 0.75 lb Al/ac.
Lorsban	1B	•••	•	•••	Tank mix with another product for broad spectrum Lep activity; For use on cole crops,
					PHI: 21 d; use top of label rates if possible.
Acephate	1B	•	••	••	Tank mix with another product for broad spectrum Lep activity; PHI: 21 d on head
					lettuce only.
Pyrethroids	3	•	•••	•••	Tank mix with another product for broad spectrum Lep activity; PHI: varies with
•					products; use high labeled rates
Radiant	5	•••	•••	•••	Stand alone Lep, leafminer, and thrips control; PHI: 1 day on lettuce; Use rates at 5-7 oz
					depending on pest spectrum.
Proclaim	6	•••	••	•••	Stand alone Lep control; use a penetrating adjuvant; PHI: 7 day on lettuce; use at rates
					above 3.6 oz; if cabbage looper present tank-mixed with a pyrethroid.
Bt (i.e. Dipel)	11B	•	• •	•	Tank mix with another product for broad spectrum Lep activity, numerous Bt products available; PHI: 0 d -good spray coverage desirable
					Tank mix with another product for broad spectrum Lep activity; PHI: 1 day; good spray
Intrepid	18A	•••	•••	••	coverage desirable; mix with a pyrethroid for best results
					Tank mix with another product for broad spectrum Lep activity; PHI: 1 day, good spray
Avaunt	22	•••	•••	••	coverage desirable, use higher rates for best control
Belt	28	•••	•••	••	Stand alone Lep control; PHI: 1 day on lettuce, Use at higher rates.
	20				Stand alone Lep and leafminer control; PHI: 1 day for lettuce- Use at or above 5 oz. for
Coragen	28	•••	•••	•••	best residual effectiveness.
Frigal	20			•••	Foliar only; Stand alone Lep, whitefly and leafminer control; PHI: 1 day for lettuce- Use
Exirel	28	•••	•••	•••	at or above 13 oz. for best residual effectiveness.
Verimark	28	•••	•••	•••	Soil only; Stand alone Lep, whitefly and leafminer control; Use at or above 10 oz. for
					best residual effectiveness.
Voliam Xpress	28+3	•••	•••	•••	Stand alone Lep and leafminer control; PHI: 1 day forlettuce; Use higher rates (8 oz or >
Voliaili Apress	2013				for best residual effectiveness.
Volium Flexi	28+4A	•••	•••	•••	Stand alone Lep and leafminer control; PHI: 7 day for lettuce; Has aphid activity. Use
					higher rates for best residual effectiveness.
Durivo	28+4A	•••	•••	•••	Soil only; Stand alone Lep and leafminer control; PHI: 30 day for lettuce; Use at 13 oz.
	20.47				for best residual effectiveness. Has aphid activity.
Vetica	28+16	•••	•••	••	Stand alone Lep control; PHI: 7 day for lettuce; Has whitefly immature activity. Use at 17
				1	oz for best residual effectiveness.

Good residual control (7-14 d)

Marginal residual control (4-6 d)

Poor residual control (1-3 d)

¹ IRAC Mode of Action - for more infor go to - http://www.irac-online.org/

^{*} always consult the label before applying any of these products