



ARIZONA AND NEW MEXICO DAIRY NEWSLETTER

COOPERATIVE EXTENSION
The University of Arizona
New Mexico State University

NOVEMBER, 2002

Evaluating Herd Milk Quality Using DHI Somatic Cell Counts

Robert R. Peters

*Professor and Extension Dairy Specialist
Department of Animal and Avian Sciences
University of Maryland*

Introduction

The somatic cell count (SCC) option is now widely selected by dairy producers in the DHI program for monitoring udder health. This is an excellent program because it provides the opportunity to evaluate the extent of subclinical mastitis in the herd and if needed, some reports may be generated to help diagnose the source of the mastitis. This is an opportunity for more profitable dairying since it is known that SCC are the single most important factor in accounting for variation in DHI herd average milk production (25). Before going any further it is important to discuss mastitis in terms of the bigger picture. Without covering some of the basic aspects of mastitis and the terminology that goes with it, understanding the management information in the DHI reports will be difficult. The points that need brief discussion to introduce this topic are the cause of mastitis, natural defense mechanism to udder infection, major forms of mastitis, and the economics of the disease. Excellent texts with an overview of mastitis are *Current Concepts of Bovine Mastitis* (2) and *Winning the Fight Against Mastitis* (20).

(continued on page 3)

Originally presented at the Arizona Dairy Production Conference, October 17, 2002, Phoenix, Arizona

New Mexico State University Extension Dairy Website:
<http://www/nmsu.edu/~dairy>

The following videos are available for checkout from Mike Looper, New Mexico State University. To obtain a video call Kathy Bustos, (505) 646-3325 or kbustos@nmsu.edu and the video will be sent in the mail, pending availability. There is only one copy of each video available, so we request that videos be returned within two weeks. Note that four of the videos contain an English and Spanish version.

1. The Milking School. Utah State University. Spanish and English. 30 minutes
2. Fitting and Showing Your Dairy Animal....A Winning Experience. Department of Dairy Science, University of Wisconsin. 20 minutes
3. Proper Milking Procedure. University of Florida. Spanish and English. 12 minutes
4. Milking Machine Maintenance. University of Florida. Spanish and English. 16 minutes
5. The Basics of Vacuum and Milking Systems. DHIA Services, 1991. 53 minutes
6. Understanding Dairy Cattle Behavior to Improve Handling and Production. Livestock Conservation Institute, 1992
7. Managing Milking/Ordenar Lecheria. Spanish and English. 1999. 33 minutes
8. Get Milk! Joining A Dairy Crew. University of New Hampshire, 1999. 45 minutes

Need to Calculate Production Costs?

University of Wisconsin dairy farm management specialist, Gary Frank, has developed a Excel spreadsheet to calculate variable cost of production and total cost of production. To access the spreadsheet, go to <http://www.wisc.edu/dairy-profit>, click on Decision Making Tools, then go to [costcwt.xls](#).

Issued in furtherance of Cooperative Extension on work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of agriculture, James A. Christenson, Director, Cooperative Extension, College of Agriculture, The University of Arizona. The University of Arizona College of Agriculture is an Equal Opportunity Employer, authorized to provide research, educational information and other services only to individuals and institutions that function without regard to sex, race, religion, color, national origin, age, Vietnam Era Veteran's status, or disability.



University of Arizona
Department of Animal Sciences
P O Box 210038, Shantz Bldg, Room 205
Tucson, AZ 85721



New Mexico State University
Extension Animal Resources
Box 30003 Dept. 3AE
Las Cruces, NM 88003

Evaluating Herd Milk Quality (continued from page 1)

Basics of Mastitis

Mastitis is an inflammation of the mammary gland. The source of that inflammation is usually microorganisms that gain access to the mammary gland through the orifice of the teat end. Microorganisms that cause disease are called pathogens. In mastitis control programs, these organisms are categorized as either contagious or environmental pathogens. Like any disease caused by bacterial infection, mastitis is a numbers game. Greater bacterial exposure translates to a greater rate of infection and more mastitis. To date, most of the progress in controlling mastitis has resulted from reducing exposure of cow's teats to pathogens, and from the widespread use of antibiotics. That is why management plays such a big part in this disease. Dr. Jim Jarrett, regular writer for the Cowside Practice column in Hoard's Dairyman, sums mastitis control up in three words: clean, dry, and comfortable. All of these are under the direct control of the manager. Controlling bacterial infection also involves the mammary gland's resistance to infection. Two examples of steps which may be taken to improve resistance to infection are supplementing the cow's diet with the micronutrients or minerals and vaccination against coliform mastitis.

The natural defense of the cow to udder infection calls the immune system into action. When the tissue in the udder senses a foreign body and the irritation that accompanies the pathogenic invasion, white blood cells from bone marrow storage pools are transported to the site of infection via the bloodstream. From the bloodstream these white blood cells migrate into the secretory tissue and in milk to seek out and kill the pathogens. The number of white blood cells which enter the mammary gland depends on the type and number of pathogens which gain access to the udder. The range is from hundreds of thousands to millions of white blood cells. The literature indicates that a SCC of 200,000 from an individual quarter or cow is the upper limit for normal milk from uninfected quarters. Thus, a 200,000 SCC is a threshold that may be used to distinguish infected cows from uninfected cows

The term SCC is used rather than a white blood cell count because it includes both epithelial cells sloughed from the milk duct of the udder and white blood cells. Approximately 15-17% of the cell count from uninfected cows are sloughed epithelial cells and the remainder are white blood cells (17). In uninfected quarters, macrophages and neutrophils each make up about 30% of the somatic cell count and lymphocytes 25%. But in quarters with subclinical and clinical mastitis a shift in cell type takes place such that approximately 90% of the cells are neutrophils (3,18). Neutrophils are the primary type of white blood cell which seeks out, engulfs, and digests pathogens.

Three forms of mastitis which are commonly discussed are subclinical, clinical, and chronic. Subclinical mastitis is not discernable with the naked eye. The milk and udder appear normal yet the milk composition has changed. The primary change which is measured is the increase in concentration of white blood cells or SCC. Cowside tests such as the California Mastitis Test may be used to detect this type of mastitis. DHI laboratories use electronic counters. Symptoms of clinical mastitis are flakes, clots, or blood in the milk, a swollen quarter, or perhaps an increased body temperature or other symptoms affecting the cow's body. Chronic mastitis is characterized as mastitis of long duration. Usually episodes of clinical mastitis appear sporadically but in the intervening periods between clinical episodes, the udder infection only subsides to the subclinical form of the disease. The DHI SCC does not estimate clinical mastitis. It is a system to determine the extent of subclinical mastitis or the development of new udder infections. Chronic mastitis may also be measured depending on the processing center and type of custom report generated.

It is estimated that the economic cost of mastitis is \$184.40 per cow per year (2). Subclinical mastitis caused the largest proportion (66%) of the cost because this form of the disease reduces milk production. Although costs associated with clinical cases such as additional culling, veterinary, and medicine costs are direct out-of-pocket expenses, together they only represent 34% of the total cost. Milk production is still the number one economic loss. A recent article in 2002 by S. L. Ott (15) reported that the average bulk tank somatic cell count (BTSCC) from 15,806 herds in 14 states. Ninety percent of the herds were located in four states (WI, OH, MI, and PA). Bulk tank somatic cell counts increased each year from 307,000 cells per ml in 1997 to 320,000 cells per ml in 2001 (Table 1). Ott and Novak (16) reported that herds with BTSCC less than 200,000 cells per ml generated \$104 per cow more in

annual value of production than herds with 200,000 to 399,999 cells per ml and \$292 per cow more than herds with 400,000 or more cells per ml.

Arizona DHI values which estimated BTSCC were similar to nationwide statistics (26). For example, the rolling 12 month average herd SCC in August, 2002, for 32 herds on the Provo DHI processing center was 323,000 cells per ml. The rolling 12-month average ranged from 180,000 to 486,000 for these 32 herds. Monthly average for August 2002 of the 22 herds on the SCC program were also reviewed. As expected in the middle of summer heat, August 2002, the average (363,000 cells per ml) and range (246,000 – 508,000) were slightly higher than the 12-month rolling average.

One cost which has been left out of the estimates by the National Mastitis Council are milk quality premiums. Usually the most limiting factor to receiving these premiums are a low SCC and thus, indirectly, managing the herd for low SCC usually has the added economic benefit of milk quality premiums. Dr. John Fetrow has developed a spreadsheet which calculates the economic cost of mastitis on a particular dairy farm (8). This spreadsheet included the premiums for milk quality and the penalty for high somatic cell counts. The attractive aspect of the spreadsheet is that it calculates the mastitis loss relative to goals. That is, it is not reasonable or profitable to eliminate all mastitis and thus all mastitis costs. Rather, there is some baseline level of clinical mastitis, culling, and somatic cell counts that are reasonable for a given farm. When current herd somatic cell count level were inputted in the spreadsheet and a goal level of 150,000 somatic cell count was set for the University of Maryland dairy herd, the major economic loss was the lost milk premiums. Our milk market, Land O Lakes, offers a milk quality program which ranges from -\$0.10 to \$0.50 per cwt. Maintaining BTSCC between 126,000 and 150,000 pays \$0.40 per cwt. While milk loss due to high SCC accounts for the largest economic loss for the majority of herds, it is not difficult to imagine that clinical mastitis, culling and death could be the major sources of economic loss in herds with excellent mastitis control as measured by somatic cell count.

Low Somatic Cell Counts and Risk of Clinical Mastitis

Some dairy farmers have been reluctant to strive for low SCC's in their herds because they fear their cows will be more susceptible to mastitis. Results of Virginia Tech research (4) suggest that animals with low initial SCC's (less than 100,000 cell per ml) in first lactation have no greater risk of increased mastitis incidences in second and subsequent lactations when compared to animals with higher initial counts. Hogan et al. (10) reported that minor pathogens increased SCC approximately 100,000 to 300,000 cell per ml compared to uninfected quarters but colonization of the udder with these organisms did not prevent infection by environmental pathogens. Matthews and Harmon (11) observed that infection by *Staph aureus*, streptococci, and coliforms also was not prevented in quarters first infected with minor pathogens. Other studies (3, 18) indicate that the SCC level must reach at least 900,000 cells per ml before a cow can reliably prevent udder infection.

A study of Erskine et al. (7) indicates that the rate of clinical mastitis is similar between low and high SCC herds but coliform mastitis is higher in low SCC herds. Eighteen dairy herds with "low" and "high" SCC were selected for six months of study. Twelve herds had 12-month DHI SCC of less than or equal to 150,000 cells per ml (low SCC) and six had a 12-month SCC average of 700,000 cells per ml or greater (high SCC). Quarter milk samples were collected for bacteriologic culture from all cows with clinical mastitis. One of the major findings of the study was that on average there was little difference in clinical mastitis between herds with low (4.23 infections/100 cows/month; range, 0.42 to 10.25 infections) and high (2.91 infections/100 cows/month; range, 1.33 to 3.92) SCC. Notice the range for the low SCC is much greater than high SCC. Thus, some herds with low SCC maintained a very low incidence of clinical mastitis perhaps suggesting superior management. The one major concern, however, was the incidence of clinical coliform mastitis was significantly higher in low SCC than high SCC herds. The authors of the study made no attempt to explain this finding from observations of herd management. But it is well known that clinical coliform mastitis is a management problem. If clinical mastitis is a problem for a producer, it appears that the effective management practices need to be implemented rather than blame the low SCC.

A more recent study (19) from England indicates that producers need to manage the transition cow very carefully. This 12-month prospective study of clinical mastitis was conducted in 482 British dairy herds with a bulk milk SCC of less than 150,000 cell per ml. Clinical mastitis averaged 23.1% of the cows per herd (range 0 to 80). Over 22% of

quarter cases occurred in the first 7 days of lactation and over 50% occurred in the first 90 days of lactation. Cluster analysis indicated that the main difference between herds with a low and average incidence of mastitis was the proportion of clinical cases occurring in the first 7 day of lactation, 14% compared with 44%, respectively. Another recent report (5) from Wisconsin indicates that summer is a vulnerable time for fresh cows. In this study of 51 herd the authors found elevations in SCC and prevalence of infection during the summer (July through September) were associated with significant increase in fresh cow and herd new infection rates. These studies point to the importance of excellent sanitation 2 weeks before and around the time of parturition when the cow is very susceptible to mastitis infection. In total, the preponderance of research evidence suggests that the economic benefits reaped from lower SCC should be the decisive factor in a mastitis control program rather than the fear of increased risk of mastitis.

Interpreting DHI Somatic Cell Counts

SCC and Milk Production. Somatic cell counts are reported on DHI SCC in three ways: either as the actual cell count per ml of milk, the weighted SCC, or the linear score (LS) Dr. George Shook, University of Wisconsin, developed and promoted the LS system because it more clearly relates milk yield losses due to subclinical mastitis than actual or raw SCC. The actual SCC is converted to LS using the range for each score in Table 2.

In Table 3, the relationships among LS, SCC and milk yield are shown. Each increase in LS is associated with a decrease in milk production of 1.5 lbs per day or 400 lb per lactation regardless of whether the score increases from 2 to 3 or 7 to 8. Notice, however, that the milk yield loss of first lactation cows is one-half that of older cows. But it should be emphasized that the average lactational LS should be used to accurately calculate milk loss. Loss estimates based on a single SCC test of individual cows are not reliable.

SCC and Infection Status. Somatic cell count has been accepted as the best quantitative index of inflammation in the bovine mammary gland. The primary factor affecting somatic cell counts is infection status. Based on a number of studies the uninfected quarters have the lowest cell counts, those infected with major pathogens have the highest cell counts and minor pathogens elicit a cellular response slightly greater than that found in uninfected quarters (Table 4).

Common Misinterpretations. There is a long list of factors other than intramammary infection that can be blamed for deteriorating udder health. Studies from the scientific literature support the idea that several factors are associated with changes in herd SCC (21). Such things as age of the cow, stage of lactation, season, stress, stray voltage, herd health, day-to-day variation, and estrus may all be related to SCC changes. But the relationship between these factors and the SCC is not cause-and-effect. The principle to remember is that these other factors may exert an effect on the SCC but only if the cow or herd also harbor intramammary infection.

Age and Stage of Lactation. On the DHI-SCC reports, the cell count LS is routinely categorized by age and stage of lactation. To aid in the interpretation of these two factors, the results of a study by Sheldrake et al. (23) are helpful. SCC remain low (100,000 to 200,000 cells per ml) regardless of age. Quarters infected with minor pathogens have SCC of 200,000 to 500,000 and in *Staph aureus* infected quarters, the SCC ranges from 500,000 to 2 million. Next to infection, lactation number has the greatest influence on SCC and consequently, a review of the older cows DHI SCC record is one good place to look for the source of a mastitis problem and for potential culls.

Within lactation, the SCC for uninfected quarters is inversely related to the lactation curve (Figure 2). The SCC is elevated at freshening, lowest at peak to mid-lactation and rises in late lactation. The amount of change in SCC of uninfected cows, however, is small. In Sheldrake's study, the SCC increased only 80,000 – from 83,000 cells per ml at 35 days postpartum to 160,000 cells per ml at 285 days postpartum. Thus, when interpreting the lactation trends on DHI-SCC report, we should expect no, or minimum, increases across a lactation in a herd with an effective mastitis control program.

Diurnal SCC Variation. Reneau (21), in his review of the SCC literature, reports that the one valid concern when interpreting DHI SCC are those dairy producers on a.m.-p.m. testing (where only one milking is collected). Studies quoted in the review indicate a diurnal variation of 2 to 5-fivefold due to time of sample collection. A problem that

dairy producers on the a.m.-p.m. DHI SCC program may encounter is the misclassification of uninfected cows since an increase over a fairly low threshold will suggest a new infection. The diagnosis of cows infected with major pathogens should not be affected by diurnal variation in SCC.

Threshold Somatic Cell Count. This term does not appear on the DHI SCC report but it is implied in the way that the data are categorized on SCC reports. The threshold SCC is a value that theoretically separates uninfected cows from the infected animals. Reneau from Minnesota, reviewed the literature and concluded that using 283,000 SCC (DHI LS of 5 or greater) as a threshold is 75 to 80% efficient in correctly classifying infected and uninfected animals (21). This criteria is used in the Provo Record Processing Center (*Table 5*).

North Carolina Dairy Records Processing Center has 5 categories on the monthly herd summary. In their literature, they stress the relationship of SCC to milk yield rather than to infection status (*Table 6*).

When reviewing the DHI SCC report it is important to remember that the threshold values used to distinguish one category from another were set to facilitate analysis and interpretation of the data. The threshold is not an infallible test that establishes the presence or absence of infection in every case. Eberhart et al. (6) reported the probabilities of a cow being infected with minor and major pathogens at a given DHI-SCC (*Table 7*).

The conclusion one can draw from the above data is that a threshold set at any of the seven SCC levels would have two types of errors associated with it: false positive and false negative results. At the low end of the SCC range, false positives would be anticipated for major pathogens and a high proportion of false negatives for minor pathogens. Conversely, at the upper end of the SCC range, few false positives would happen for major pathogens or false negatives for minor pathogens. Therefore, it becomes apparent that on the basis of infection probability, there is no perfect threshold level. The rationale behind choosing the thresholds in the DHI-SCC program is that they are levels that are useful in monitoring a dairyman's mastitis control procedures. Other purposes for setting threshold SCC could be culling, culturing, and treatment (i.e. selective dry cow therapy). And each of these purposes could have different thresholds depending on the amount of confidence desired in making a correct choice.

Prevalence. This is the number of existing infections and their duration. It is another term that does not appear on the DHI SCC report but is implied depending on the source of the records. Prevalence, or level of mastitis, is a relatively static measure of herd mastitis and because it changes slowly, may not reflect recent management changes that the farmer has made. Infections of contagious organisms last weeks or months and consequently, prevalence is more indicative of contagious than environmental mastitis. Prevalence of environmental streptococci and coliform rarely exceed 10 and 5%, respectively.

Goals for prevalence can be presented as % Low (Provo records) or in LS. Suggested goals are stated in (*Table 8*).

New Infection Rate. Improved management or deteriorating udder health should be immediately reflected in this parameter. Sixty percent of environmental streptococci and 69% of coliform infections last less than 30 days. Thus, new infections are probably more important for environmental than contagious mastitis since duration of infection is short. Provo records present this information on the monthly herd report on the somatic cell summary section near the upper left hand corner. Under the column abbreviated CHG. 5, the two sub-headings are % UP and % Down for lactation number 1, 2, 3+ and Total. The new infection rate would be indicated in the % UP column. Any significant rise in this category would indicate that the new infection rate is increasing.

If advanced DHI computer programs with graphic capabilities are available, more in-depth analyses may be conducted. Trend graphs can provide producers and their advisors with a picture of what is occurring in their herds. Graphing linear score versus days in milk (by lactation group) provides a detailed picture of where, in the production cycle, infections are occurring

Scatter graphs may be used to graphing test day linear score (LS) by previous linear score (PLS) clearly illustrates the distribution of animals in one of four groups. Those groups include non-infected animals (LS < 4.0 and PLS < 4); newly infected animals (LS > 4.0 and PLS < 4.0); cured animals (LS < 4.0 and PLS > 4.0); and chronic animals (both LS and PLS > 4.0). This graph can be examined over time to determine whether performance is getting better or worse? If a large number of chronic cows exist within the herd, look for an infectious organism. If new

infections are high, as well as the number of cured animals, examine environmental conditions and milking procedures.

Diagnostic Uses of DHI SCC Using Paper Records

The two general types of paper records that should be reviewed to determine the status of the herd's udder health are the Monthly Herd Summary and Individual Cow Records. These may be separate reports or combined into one depending on the record processing center. The Herd Summary should be used to evaluate the overall management such as heifer, dry cow and lactation management. Once these areas have been reviewed, more specific decisions can be made from the individual cow records.

Probably the most important point to remember when attempting to make diagnoses from DHI-SCC is that one needs to study the trends over time. A single SCC on one cow or even a single herd summary for one month on the herd has limited value. But the trends that emerge from several months of DHI-SCC testing are meaningful and useful diagnoses can be made from them.

Monthly Herd Summary

Overall Management. The annual summary should indicate the prevalence of mastitis or economic loss during the year and whether the trend in mastitis control is maintaining current progress, improving or worsening over time. If mastitis is more prevalent during certain seasons of the year, this also should be evident from reviewing these results. Further study of the current monthly data are helpful in evaluating more specific areas of management.

Heifer Management. One very important concept in mastitis control is that overall udder health can be no better than the current status of the heifers. Dairy producers must excel in this area to maintain an excellent udder health program. The prevalence should be extremely low as indicated by the percent low or average LS. In addition, the first LS average in the stage of lactation summary should be low and remain low for the duration of lactation. If the first test consistently is above 3.0, then calf housing and raising practices and the maternity care for these animals needs evaluation.

Dry Cow Management. This area of management can be evaluated using the first test for stage of lactation. If the average LS consistently exceeds the monthly average for the herd or is as high as the last stage of lactation LS, dry cow management may need attention. Such practices as drying off strategy, administration of dry cow therapy, housing and environment, maternity care, and number of chronic mastitis cows could be at fault.

Lactation Management. The trends in LS for each stage of lactation and lactation number should be evaluated to determine whether milking management might be the area needing management changes. Since most first lactation cows are usually uninfected, they will be most sensitive to deficiencies in milking management, particularly if contagious mastitis is involved. If new infections are developing during lactation in this group, these will be reflected with higher LS (higher than first stage of lactation LS), particularly in mid lactation. The exact amount of change in LS during lactation is difficult to define. Another important point to consider are the number of cows represented in each category and stage of lactation. Averages of small numbers can obviously be misleading. It also would be helpful to review several of the last herd summaries to increase your confidence when drawing conclusions.

Individual Cow Data

Potential uses of individual cow records are culling, early dry-off, segregation and withholding of milk from high SCC to ensure payment of milk quality premium.

Culling. Candidates for culling would be those that run consistently high (LS of 6 or greater) and do not respond to dry cow therapy. *Staph aureus* cows in this category should be at the top of the cull list.

Early dry-off. Cows that are consistently running a high SCC and have a high potential for milk production or breeding stock may merit early dry-off. The objective is similar to that of culling since this practice removes the source of infection and high SCC from the lactating herd. This option probably has limited value for most dairy producers.

Segregation. My experience in Maryland is that dairy producers will not, as a rule, segregate cows (flat barn or parlor herds) on the basis of DHI SCC. The more practical alternative has been to visually identify those cows with a legband and backflush the milker with a sanitizer solution.

Withholding Milk. An individual cow can account for a significant percentage of the total bulk tank SCC. Management reports can be ordered to determine what contribution high SCC cows are making to the BTSCC. Usually it is not economical to cull a cow due to high SCC because the value of her milk is greater than premiums gained by removing the high SCC cow. But it will usually be a good decision to cull very high SCC cows if replacements are available.

Use of Individual DHI SCC for Treatment

Lactational antibiotic treatment based only on high SCC is not recommended and will yield disappointing results. Research at Cornell University (12), University of Wisconsin (27), and Virginia Tech (22) has demonstrated that this practice *does not* increase milk production or eliminate udder infections which cause the high SCC.

SCC Not a "Stand Alone" Tool

The DHI SCC are not a stand alone diagnostic procedure for evaluating udder health. It is, however, a good place to start in when reviewing the effectiveness of the dairy producer's mastitis control practices. But realistically, if the DHI-SCC results do indicate potential problems, then data from other procedures such as bulk tank or individual cow cultures, milking machine evaluations and observations from a herd visit must be included before specific conclusions and recommendations can be made for an individual herd.

Two examples in the literature illustrate why data other than SCC are required to diagnose mastitis problems. Hoblet et al. (6) made a presumptive herd diagnosis of environmental mastitis on the basis of observed nature of clinical mastitis and consistent history of low DHI SCC (87% of cows <283,000) and bulk tank SCC (annual average 137,000). Yet results of their study indicated that coagulase-positive Staphylococci, including *Staph aureus*, were repeatedly cultured from cows with SCC less than 200,000 cells per ml. Thus, bacteriologic culture was necessary to accurately diagnose the source of the mastitis even though SCC profiles suggested contagious mastitis was unlikely. Another example is the finding of McDonald reported at the 1989 Annual National Mastitis Council Meeting (10). His experience indicates that in 12 large Washington herds that were monitored over a two-year period that one-half of *Staph aureus* infected cows had individual SCC of less than 300,000. He suggests that culling based on DHI SCC would be less than ideal and that bacterial culture is a wise practice, especially with the advent of the new Pro-Staph antibody test available through DHI, before making any culling decisions based on mastitis.

References

1. Andrews, R.J., B.J. Kitchen, W.S. Kwee, and F. Duncalfe. 1983. Relationship between individual cow somatic cell counts and the mastitis infection status of the udder. *Aust. J. Dairy Technol.* 38:71.
2. Anonymous. 1996. Current concepts of bovine mastitis. Natl Mastitis Council, 4th Edition. Madison, WI.
3. Bright, S.A., W.D. Schultze, M.J. Paape, and R.R. Peters. 1987. Effect of intramammary devices on the outcome of induced *Escherichia coli* infection of bovine mammary quarters. *Am. J. Vet. Res.* 48:1290.
4. Coffey, E.M., W.E. Vinson, and R.E. Pearson. 1986. Somatic cell counts and infection rates for cows of varying somatic cell count in initial test of first lactation. *J. Dairy Sci.* 69:552.
5. Cook, N. B., T. B. Bennett, K.M. Emery, and K. V. Nordland. 2002. Monitoring nonlactating cow intramammary infection dynamics using DHI somatic cell count data. *J. Dairy Sci.* 85: 1119-1126.
6. Eberhart, R.J., H.C. Gilmore, L.J. Hutchinson, and S.B. Spencer. 1979. Somatic cell counts in DHI samples. Page 32 in 18th Proceedings of Annu. Mtg. Of Natl. Mastitis Council, Louisville, Ky.
7. Erskine, R.J., R.J. Eberhart, L.J. Hutchinson, S.B. Spencer, and M.A. Campbell. 1988. Incidence and types of clinical mastitis in dairy herds with high and low somatic cell counts. *J. Am. Vet. Med. Assoc.* 192:761.
8. Fetrow, J., S. Stewart, S. Eicker, R. Farnsworth, and R. Bey. 2000. Mastitis: an economic consideration. Pages 3-47 in the 39th Ann Mtg of the Natl. Mastitis Council, Atlanta, GA.
9. Hoblet, K.H., J.S. Bailey, and D.E. Pritchard. 1988. Coagulase-positive staphylococcal mastitis in a herd with low somatic cell counts. *J. Am. Vet. Med. Assoc.* 192:777.
10. Hogan, J.S., K.L. Smith, D.A. Todhunter, and P.S. Schoenberger. 1988. Rate of environmental mastitis in quarters infected with *Corynebacterium bovis* and staphylococcus species. *J. Dairy Sci.* 71:2520.
11. Matthews, K.R. and R.J. Harmon. 1989. Effect of naturally occurring coagulase-negative staphylococci infections on new infections by minor and major pathogens in cattle. *J. Anim. Sci.* 67(Suppl 1)/*J. Dairy Sci.* 72(Suppl 1):21(Abstr.).
12. McDermott, M.P., H.N. Erb, R.P. Natzke, F.D. Barnes, and D. Bray. 1983. Cost benefit analysis of lactation therapy with somatic cell counts as indications for treatment. *J. Dairy Sci.* 66:1198.
13. McDonald, J.S. 1989. Investigation of and resolution of herd mastitis due to *Staphylococcus aureus*. Page 89 in 28th Proceedings of Annu. Mtg. of Natl. Mastitis Council, Tampa, FL.
14. Natzke, R.P., R.W. Everett, and D.S. Postle. 1972. Normal milk somatic cell counts. *J. Milk Food Technol.* 35:261.
15. Ott, S. L., 2002. Five year trend shows rise in U.S. bulk tank somatic cell counts. *Udder Topics, Natl Mastitis Council Newsletter*: 25 (No. 3) p. 1.
16. Ott, S. L., and P. R. Novak. 2001. Association of herd productivity and bulk tank somatic cell counts in U.S. dairy herds in 1996. *J. Am. Vet. Med. Assoc.* 218 (8) 1325-1330.
17. Paape, M.J. and R.H. Miller. 1988. Identification of epithelial cells in bovine milk. *J. Dairy Sci.* 71(Suppl 1):232(Abstr.).

18. Paape, M.J., W.D. Schultze, A.J. Guidry, W.M. Kortum, and B.T. Weinland. 1981. Effect of an intramammary polyethylene device on the concentration of leukocytes and immunoglobulins in milk and on the leukocyte response of *Escherichia coli* endotoxin and challenge exposure with *staphylococcus aureus*. Am. J. Vet. Res. 42:774.
19. Peeler, E. J., M.J. Green, J.L. Fetzpatrick, and L.E. Green. 2002. Study of clinical mastitis in British dairy herds with bulk tank somatic cell counts less than 150,000 cells/ml. Vet Rec. 11 (6):170-176.
20. Philpot, W. N., and S.C. Nickerson. 2000. Winning the fight against mastitis. Published by Westfalia-Surge, Inc. Naperville, IL.
21. Reneau, J.K. 1986. Effective use of dairy herd improvement somatic cell counts in mastitis control. J. Dairy Sci. 69:1708.
22. Seymour, E.H., G.M. Jones, and M.L. McGilliard. 1989. Effectiveness of intramammary antibiotic therapy based on somatic cell count. J. Dairy Sci. 72:1057.
23. Sheldrake, R.F., R.J.T. Hoare, and G.D. McGregor. 1983. Lactation stage, parity, and infection affecting somatic cells, electrical conductivity, and serum albumin in milk. J. Dairy Sci. 66:542.
24. Schultz, L.H. 1977. Somatic cells in milk physiological aspects and relationship to amount and composition of milk. J. Food Prot. 40:125.
25. Shutz, M.M., R.D. Appleman, G.R. Steurnagel, and L.B. Hansen. 1985. Determining and assessing management traits that influence herd averages for milk production. J. Dairy Sci. 68(Suppl 1):326(Abstr.).
26. Smith, S. 2002. Personal communication. DHI Provo, Provo, Utah.
27. Timms, L.L. and L.H. Schultz. 1984. Mastitis therapy for cows with elevated somatic cell counts or clinical mastitis. J. Dairy Sci. 67:367.

Table 1. Trends in Bulk Tank Somatic Cell Counts
(BT SCC) Among 15,806 Herds, 1997-2001.

<u>Year</u>	<u>BT SCC</u> <u>1000's</u>
1997	307
1998	310
1999	312
2000	315
2001	320

Reference: S.L. Ott (15)

Table 2. SCC that correspond to linear score

<u>Linear Score</u>	<u>Cell Counts (X10³)</u>	
	<u>Midpoint</u>	<u>Range</u>
0	12.5	0-17
1	25	18-35
2	50	35-70
3	100	71-140
4	200	141-282
5	400	283-565
6	800	566-1,130
7	1,600	1,131-2,262
8	3,200	2,263-4,525
9	6,400	4,526-

Table 3. Milk loss estimates from linear score

<u>Linear Score</u>	<u>Mid-point SCC</u>	<u>Milk Yield Loss</u>		
		<u>lb/day¹</u>	<u>lb/lactation</u>	
			<u>First Calf</u>	<u>Older Cows</u>
2	50	0	0	0
3	100	1.5	200	400
4	200	3.0	400	800
5	400	4.5	600	1,200
6	800	6.0	800	1,600
7	1,600	7.5	1,000	2,000
8	3,200	9.0	1,200	2,400

¹Losses in first lactation are one-half these amounts.

Table 4. Summary of Mean SCC or SCC Range by infection status among several studies

Reference	Mean SCC or SCC Range		
	Uninfected	Minor	Major
Schultz, 1977 (19)	170,000	227,000	998,000
Eberhart et al., 1979 (4)	165,000	364,000	1,061,000
Natzke et al., 1972 (11)	214,000	...	504,000-1,470,000
Sheldrake et al., 1983 (18)	100,000-175,000	200,000-500,000	500,000+
Andrew et al., 1983 (1)	147,000	...	556,000

Reference: J.K. Reneau (21)

Table 5. DHI-Provo Threshold Standard's for Infection Status

<u>SCC Category</u>	<u>Linear Score</u>	<u>Interpretation</u>
Low	0-4	Noninfected
Medium	5-6	Infected
High	7-9	Infected

Table 6. North Carolina Threshold Standards.

% of cows with Linear Score (LS) in each of five categories

LS	0, 1, 2, 3
LS	4
LS	5
LS	6
LS	7, 8, 9

Table 7. Probability of infection with minor and major pathogen at various concentrations of somatic cell counts.

<u>SCC X10³</u>	<u>Probability of Minor Pathogen</u>	<u>Probability of Major Pathogen</u>
<100	.48	.06
100-200	.59	.17
200-300	.50	.34
300-400	.44	.45
400-500	.35	.51
500-600	.28	.65
>600	.19	.79

Reference: Eberhart et al. (6).

Table 8. Goals for % Low and Linear Score on Provo DHI Records.

<u>% Low (Linear Score 0-4)</u>	<u>Linear Score</u>	<u>Rating</u>
≥90	1.0-2.5	Excellent
80-89	2.6-3.2	Good
70-79	3.3-3.8	Average
60-69	3.9-4.4	Below Average
<60	≥4.5	Poor

HIGH COW REPORT OCTOBER, 2002

MILK

Arizona Owner	Barn #	Age	Milk	New Mexico Owner	Barn #	Age	Milk
* Stotz Dairy West	13439	4-1	41,080	Ken Miller Dairy	625	5-01	43,611
* Stotz Dairy West	19194	6-2	38,950	* Providence Dairy	3265	5-01	39,960
* Stotz Dairy West	6764	6-5	38,080	* Hide-Away Dairy	3377	4-03	39,460
* Stotz Dairy West	13280	4-1	37,290	Mariposa Dairy Farm	4229	5-05	37,980
* Stotz Dairy West	19633	4-5	37,040	* Hafliger Dairy	3750	7-06	37,840
* Stotz Dairy West	13321	4-2	36,980	* Do-Rene Dairy	1461	5-06	37,440
* Dutch View Dairy	320	5-6	36,790	S.A.S. Dairy	1664	7-00	37,372
* Saddle Mountain Dairy	248	2-11	36,610	Pareo Dairy	1053	7-07	37,154
* Stotz Dairy West	11471	5-6	36,610	Mariposa Dairy Farm	MF396	2-00	36,510
* Stotz Dairy West	5545	8-6	36,440	McCatharn Dairy	3009	6-03	36,301

FAT

* Stotz Dairy West	19194	6-2	1751	* Hafliger Dairy	3750	7-06	1454
* Stotz Dairy West	19633	4-5	1491	Mariposa Dairy Farm	4229	5-05	1428
* Stotz Dairy West	13334	4-1	1424	New Direction Dairy	1514	2-02	1395
* Stotz Dairy West	13537	3-11	1423	Pareo Dairy	8491	5-09	1366
* Stotz Dairy West	10411	8-5	1402	Mariposa Dairy Farm	MF396	2-00	1360
* Stotz Dairy West	13439	4-1	1387	Pareo Dairy	1400	7-01	1347
* Stotz Dairy West	19586	4-6	1361	* Hafliger Dairy	6906	4-03	1339
* Dutch View Dairy	320	5-6	1360	New Direction Dairy	1488	3-01	1321
* Stotz Dairy West	11471	5-6	1359	Pareo Dairy	19	5-01	1317
Wigwam Dairy	1647	3-7	1357	Mariposa Dairy Farm	4978	4-09	1304
				Mariposa Dairy Farm	3442	8-03	1304

PROTEIN

* Stotz Dairy West	19194	6-2	1153	Ken Miller Dairy	625	5-01	1190
* Stotz Dairy West	13439	4-1	1104	Mariposa Dairy Farm	4229	5-05	1190
* Stotz Dairy West	14801	3-0	1101	Mariposa Dairy Farm	MF396	2-00	1138
* Dutch View Dairy	320	5-6	1072	* Hafliger Dairy	3750	7-06	1132
* Stotz Dairy West	6764	6-5	1061	* Providence Dairy	3265	5-01	1128
* Stotz Dairy West	19571	4-5	1060	* Hide-Away Dairy	1644	8-06	1126
* Stotz Dairy West	19480	4-5	1057	* High Plains Dairy	72	5-09	1111
* Stotz Dairy West	13681	3-10	1057	Mariposa Dairy Farm	3442	8-03	1104
* Stotz Dairy West	11717	5-3	1053	S.A.S. Dairy	4041	4-02	1103
* Stotz Dairy West	13220	4-2	1048	Mariposa Dairy Farm	4978	4-09	1099

* 3X milking

ARIZONA – TOP 50% FOR F.C.M. ^b
OCTOBER, 2002

OWNERS NAME	Number of Cows	MILK	FAT	3.5 FCM	C.I.
* Stotz Dairy	2141	28,364	1009	28,627	14.9
* Red River Dairy	4494	27,675	984	27,899	13.8
University of Arizona Holsteins	177	26,341	945	26,716	13.8
Martha Linda Dairy	1956	25,536	935	26,206	13.4
* Mike Pylman Dairy	2700	25,179	888	25,288	14.2
* Stotz Dairy East	1355	25,105	879	25,110	12.1
* Arizona Dairy North	2643	24,691	872	24,818	14.2
* Hillcrest Dairy	2320	24,750	849	24,471	13.9
University of Arizona Brown Swiss	109	22,796	900	24,453	14.5
* Arizona Dairy South	3356	24,463	850	24,363	14.0
* Del Rio Holsteins	1279	23,623	845	23,919	13.2
Paul Rovey Dairy	425	23,813	840	23,919	13.2
Desert Ridge Dairy LLC2	446	23,669	831	23,711	13.8
* Zimmerman Dairy	1212	22,986	825	23,318	14.8
Butler Dairy	623	23,130	813	23,286	14.4
* Wigwam Dairy	1518	22,798	818	23,123	15.0
* Del Rio Brown Swiss	190	21,126	789	22,925	12.9
* Saddle Mountain Dairy	2166	23,479	769	22,622	13.3
Desert Ridge Dairy LLC	1395	21,884	804	22,502	12.6
* Danzeisen Dairy, LLC	1188	22,006	794	22,392	13.2
* Dutch View Dairy	1649	22,266	778	22,244	14.1
Gladtime West Holsteins	362	21,985	753	21,718	15.3
Treger Holsteins, Inc.	512	21,236	766	21,605	13.8
* Dairyland Milk Company	2533	21,959	742	21,527	13.7
Parker Dairy	4379	20,566	773	21,429	14.6
* Goldman Dairy	2012	21,183	748	21,290	13.7
OWNERS NAME	Number of Cows	MILK	FAT	3.5 FCM	C.I.
* Pareo Dairy #1	1369	26,464	934	26,591	13.7
* Hafliger Dairy	1717	25,968	954	26,701	14.0
* Do-Rene Dairy	2867	25,634	891	25,534	13.7
McCatharn North Dairy	1056	25,062	844	24,525	13.6
* Hide-Away Dairy	2196	24,631	846	24,371	13.1
Ken Miller Dairy	386	24,330	812	23,689	14.0
* Tallmon Dairy	538	24,174	852	24,271	14.8
Pareo Dairy #2	2842	23,889	887	24,715	13.2
S.A.S. Dairy	2065	23,794	858	24,204	13.6
Price's Roswell Farm	2716	23,081	827	23,393	13.7
* Providence Dairy	2535	22,403	753	21,899	13.6
* Vaz Dairy	1567	22,335	765	22,064	14.1
* High Plains Dairy	1786	22,300	821	22,957	14.2

* 3X a day milking

^b average milk and fat figure may be different from monthly herd summary; figures used are last day/month

**ARIZONA AND NEW MEXICO HERD IMPROVEMENT SUMMARY
FOR OFFICIAL HERDS TESTED OCTOBER, 2002**

		ARIZONA	NEW MEXICO
1.	Number of herds	52	26
2.	Total cows in herd	78,883	46,106
3.	Average herd size	1516	1773
4.	Percent days in milk	86	86
5.	Average days in milk	200	202
6.	Average milk – all cows per day	56.4	57.9
7.	Average percent fat – all cows	3.6	3.7
8.	Total cows in milk	67,839	39,906
9.	Average daily milk for milking cows	66.2	67.2
10.	Average days in milk – 1 st breeding	82	72
11.	Average days open	155	148
12.	Average calving interval	13.9	13.9
13.	Percent somatic cell – linear 0-4	81	77
14.	Percent somatic cell – linear 5-6	9	16
15.	Percent somatic cell – linear 7 & above	10	6
16.	Average previous days dry	63	67
17.	Percent cows leaving herd	33	34.8
		STATE AVERAGE	
	MILK	22,102	22,167
	Percent butterfat	3.6	3.6
	Percent Protein	3.0	3.0
	Pounds fat	801	788
	Pounds protein	663	658

ARIZONA COOPERATIVE EXTENSION
U.S. DEPARTMENT OF AGRICULTURE
The University of Arizona
Tucson, Arizona 85721

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE, \$300

PRESRT STD
POSTAGE & FEES PAID
USDA
PERMIT NO G-268



THE UNIVERSITY OF
ARIZONA®

COOPERATIVE EXTENSION

Department of Animal Sciences

P O Box 21 0038

Tucson, AZ 85721 -0038

Phone: 520-621 -3375

Fax: 520-621 -9435

Email: aretig@ag.arizona.edu

Mark your calendar

- Southwest Nutrition and Management Conference, Phoenix, Arizona, February 20-21, 2003
- Arizona Dairy Day, Phoenix, Arizona, March 7, 2003
- Western Dairy Management Conference, Reno, Nevada, March 12-14, 2003